arXiv:1006.5158v2 [math.CA] 19 Jul 2010

JACOB'S LADDERS AND THE NONLOCAL INTERACTION OF THE FUNCTION $Z(t)$ WITH THE FUNCTION $\tilde{Z}^2(t)$ ON THE DISTANCE $\sim (1-c)\pi(t)$ FOR A COLLECTION OF DISCONNECTED SETS

JAN MOSER

ABSTRACT. It is shown in this paper that there is a fine correlation of the third order between the values of the functions $Z[\varphi_1(t)]$ and $\tilde{Z}^2(t)$ which corresponds to two collections of disconnected sets. The corresponding new asymptotic formula cannot be obtained within known theories of Balasubramanian, Heath-Brown and Ivic.

1. RESULT

1.1. Let (see [2], (3), (4))

$$G_1(x)=G_1(x;T,H)=\bigcup_{T\le t_{2\nu}\le T+H}\left\{t:\ t_{2\nu}(-x)<t<t_{2\nu}(x),\ 0<x\le\frac{\pi}{2}\right\}$$

$$G_2(y)=G_2(y;T,H)= \tag{1.1}$$

$$=\bigcup_{T\le t_{2\nu+1}\le T+H}\left\{t:\ t_{2\nu+1}(-y)<t<t_{2\nu+1}(y),\ 0<y\le\frac{\pi}{2}\right\}$$

$$H=T^{1/6+2\epsilon}, \tag{1.2}$$

and the collection of the sequences $\{t_\nu(\tau)\},\ \tau\in[-\pi,\pi],\ \nu=1,2,\dots$ be defined by the equation (see [2], (1))

$$\vartheta[t_\nu(\tau)]=\pi\nu+\tau;\ t_\nu(0)=t_\nu. \tag{1.3}$$

1.2. In this paper we obtain some new properties of the signal

$$Z(t)=e^{i\vartheta(t)}\zeta\left(\frac{1}{2}+it\right)$$

generated by the Riemann zeta-function. Namely, let

$$G_1(x)=\varphi_1[\mathring{G}_1(x)],\ G_2(y)=\varphi_1[\mathring{G}_2(y)] \tag{1.4}$$

where $y=\varphi_1(T),\ T\ge T_0[\varphi_1]$ is the Jacob's ladder. The following theorem holds true.

Theorem.

$$\int_{\mathring{G}_1(x)}Z[\varphi_1(t)]\tilde{Z}^2(t)\mathrm{d}t=\frac{2}{\pi}H\sin x+\mathcal{O}(T^{1/6+\epsilon}),$$

$$\int_{\mathring{G}_2(y)}Z[\varphi_1(t)]\tilde{Z}^2(t)\mathrm{d}t=-\frac{2}{\pi}H\sin y+\mathcal{O}(T^{1/6+\epsilon}), \tag{1.5}$$

Key words and phrases. Riemann zeta-function.

where

$$t-\varphi_1(t)\sim(1-c)\pi(t),\qquad t\to\infty, \tag{1.6}$$

and c is the Euler's constant and $\pi(t)$ is the prime-counting function.

Let us remind another representation of the signal $Z(t)$ given by the Riemann-Siegel formula (in its local form)

$$Z(t)=2\sum_{n<P_0}\frac{1}{\sqrt{n}}\cos\{\vartheta(t)-t\ln n\}+\mathcal{O}(T^{-1/4}),\ t\in[T,T+H],\ P_0=\sqrt{\frac{T}{2\pi}},$$

i.e. in the form of the resultant oscillation of the system of nonlinear oscillators

$$\frac{2}{\sqrt{n}}\cos\{\vartheta(t)-t\ln n\};\ \vartheta(t)=\frac{t}{2}\ln\frac{t}{2\pi}-\frac{t}{2}-\frac{\pi}{8}+\mathcal{O}\left(\frac{1}{t}\right)$$

(at the same time see, for example, the expression $Z[t_\nu(\tau)]$ by (1.3)).

1.3. Let (comp. (1.4))

$$T=\varphi_1(\overset{\circ}{T}),\ T+H=\varphi_1(\overset{\circ}{\widehat{T+H}}). \tag{1.7}$$

Since (see (1.6), (1.7))

$$\overset{\circ}{T}-\varphi_1(\overset{\circ}{T})\sim(1-c)\frac{\overset{\circ}{T}}{\ln\overset{\circ}{T}}\ \Rightarrow\ \overset{\circ}{T}-T\sim(1-c)\frac{\overset{\circ}{T}}{\ln\overset{\circ}{T}}\ \Rightarrow\ \overset{\circ}{T}\sim T$$

we have seen (see (1.2))

$$\overset{\circ}{T}-(T+H)\sim(1-c)\frac{\overset{\circ}{T}}{\ln\overset{\circ}{T}}-H\sim(1-c)\frac{\overset{\circ}{T}}{\ln\overset{\circ}{T}},$$

i.e. $\overset{\circ}{T}>T+H$. Then we have

$$\begin{aligned}&[T,T+H]\cap[\overset{\circ}{T},\overset{\circ}{\widehat{T+H}}]=\emptyset;\ T+H<\overset{\circ}{T},\\ &\rho\{[T,T+H];[\overset{\circ}{T},\overset{\circ}{\widehat{T+H}}]\}\sim(1-c)\pi(T),\end{aligned} \tag{1.8}$$

where ρ stands for the distance of the corresponding segments (comp. [5], (1.3), (1.6)).

Remark 1. Some nonlocal interaction of the functions $Z[\varphi_1(t)]$ and $\tilde{Z}^2(t)$ is expressed by the formula (1.5) where (see (1.7))

$$t\in\overset{\circ}{G}_1(x)\cup\overset{\circ}{G}_2(y)\cap[\overset{\circ}{T},\overset{\circ}{\widehat{T+H}}]\ \Rightarrow\ \varphi_1(t)\in G_1(x)\cup G_2(y)\cap[T,T+H].$$

Such an interaction is connected with two collections of disconnected sets unboundedly receding each from other (see (1.8), $\rho\to\infty$ as $T\to\infty$) - like mutually receding galaxies (the Hubble law). Compare this remark with the Remark 3 in [5].

Remark 2. The asymptotic formulae (1.5) cannot be obtained by methods of Balasubramanian, Heath-Brown and Ivic (comp. [1]).

This paper is a continuation of the series [3]-[15].

2. THE FIRST COROLLARIES

First of all, we obtain from (1.5)

Corollary 1.

(2.1)

$$\int_{\mathring{G}_1(x)\cup\mathring{G}_2(y)} Z[\varphi_1(t)]\tilde{Z}^2(t)\mathrm{d}t = \left\{ \begin{array}{lll} \frac{2}{\pi}(\sin x - \sin y)H + \mathcal{O}(T^{1/6+\epsilon}) & , & x \neq y \\ \mathcal{O}(T^{1/6+\epsilon}) & , & x = y, \end{array} \right.$$

$$\int_{\mathring{G}_1(x)} Z[\varphi_2(t)]\tilde{Z}^2(t)\mathrm{d}t - \int_{\mathring{G}_2(y)} Z[\varphi_2(t)]\tilde{Z}^2(t)\mathrm{d}t = \frac{2}{\pi}(\sin x + \sin y)H + \mathcal{O}(T^{1/6+\epsilon}).$$

Next, in the case $x = y = \frac{\pi}{2}$ we obtain

Corollary 2.

$$\int_{\mathring{G}_1\left(\frac{\pi}{2}\right)\cup\mathring{G}_2\left(\frac{\pi}{2}\right)} Z[\varphi_1(t)]\tilde{Z}^2(t)\mathrm{d}t = \mathcal{O}(T^{1/6+\epsilon}),$$

$$\int_{\mathring{G}_1\left(\frac{\pi}{2}\right)} Z[\varphi_1(t)]\tilde{Z}^2(t)\mathrm{d}t - \int_{\mathring{G}_2\left(\frac{\pi}{2}\right)} Z[\varphi_1(t)]\tilde{Z}^2(t)\mathrm{d}t = \frac{4}{\pi}H + \mathcal{O}(T^{1/6+\epsilon}),$$

where $[\mathring{T}, \overset{\circ}{\widehat{T+H}}] \subset \mathring{G}_1\left(\frac{\pi}{2}\right) \cup \mathring{G}_2\left(\frac{\pi}{2}\right)$.

3. LAW OF THE ASYMPTOTIC EQUALITY OF THE AREAS OF THE POSITIVE AND NEGATIVE PARTS OF THE GRAPH OF THE FUNCTION $Z[\varphi_1(t)]\tilde{Z}^2(t)$

Let

$$\begin{aligned} \mathring{G}_1^+(x) &= \left\{t:\ t \in \mathring{G}_1(x),\ Z[\varphi_1(t)]\tilde{Z}^2(t) > 0\right\}, \\ \mathring{G}_1^-(x) &= \left\{t:\ t \in \mathring{G}_1(x),\ Z[\varphi_1(t)]\tilde{Z}^2(t) < 0\right\}, \\ \mathring{G}_2^+(x) &= \left\{t:\ t \in \mathring{G}_2(x),\ Z[\varphi_1(t)]\tilde{Z}^2(t) > 0\right\}, \\ \mathring{G}_2^-(x) &= \left\{t:\ t \in \mathring{G}_2(x),\ Z[\varphi_1(t)]\tilde{Z}^2(t) < 0\right\}. \end{aligned} \tag{3.1}$$

Then we obtain from (2.1) by (1.1), (3.1) the following

Corollary 3.

$$\int_{\mathring{G}_1^+(x)\cup\mathring{G}_2^+(x)} Z[\varphi_1(t)]\tilde{Z}^2(t)\mathrm{d}t \sim - \int_{\mathring{G}_1^-(x)\cup\mathring{G}_2^-(x)} Z[\varphi_1(t)]\tilde{Z}^2(t)\mathrm{d}t, \quad T \to \infty. \tag{3.2}$$

Indeed, from (1.5) by (3.1) we have

$$0 < (1-\epsilon)\frac{2}{\pi}H\sin x < \int_{\mathring{G}_1(x)} \leq \int_{\mathring{G}_1^+(x)} \leq \int_{\mathring{G}_1^+(x)\cup\mathring{G}_2^+(x)},$$

and similarly

$$0 < (1-\epsilon)\frac{2}{\pi}H\sin x < -\int_{\overset{\circ}{G}{}_1^-(x)\cup\overset{\circ}{G}{}_2^-(x)} .$$

Hence, from (2.1), $x=y$, we get

$$\int_{\overset{\circ}{G}{}_1^+(x)}+\int_{\overset{\circ}{G}{}_1^-(x)}+\int_{\overset{\circ}{G}{}_2^+(x)}+\int_{\overset{\circ}{G}{}_2^-(x)}=\mathcal{O}(T^{1/6+\epsilon})=o(H)$$

(see (1.2), i.e. (3.2)).

Remark 3. The formula (3.2) represents the law of the asymptotic equality of the areas (measures) of the figures which correspond to the positive part and negative part, respectively, of the graph of the function

$$Z[\varphi_1(t)]\tilde{Z}^2(t),\ t\in\overset{\circ}{G}_1(x)\cup\overset{\circ}{G}_2(x) \tag{3.3}$$

with respect to the disconnected sets $\overset{\circ}{G}{}_1^+(x)\cup\overset{\circ}{G}{}_2^+(x),\ \overset{\circ}{G}{}_1^-(x)\cup\overset{\circ}{G}{}_2^-(x)$. This is one of the laws governing *chaotic* behavior of the positive and negative values of the function (3.3).

4. PROOF OF THE THEOREM

4.1. Let is remind that

$$\tilde{Z}^2(t)=\frac{\mathrm{d}\varphi_1(t)}{\mathrm{d}t},\ \varphi_1(t)=\frac{1}{2}\varphi(t)$$

where

$$\tilde{Z}^2(t)=\frac{Z^2(t)}{2\Phi'_\varphi[\varphi(t)]}=\frac{Z^2(t)}{\left\{1+\mathcal{O}\left(\frac{\ln\ln t}{\ln t}\right)\right\}\ln t}$$

(see [3], (3.9); [5], (1.3); [9], (1.1), (3.1), (3.2)). Thus, the following lemma holds true (see [8], (2.5); [9], (3.3)).

Lemma. For every integrable function (in the Lebesgue sense) $f(x),\ x\in[\varphi_1(T),\varphi_1(T+U)]$ the following is true

$$\int_T^{T+U} f[\varphi_1(t)]\tilde{Z}^2(t)\mathrm{d}t=\int_{\varphi_1(T)}^{\varphi_1(T+U)} f(x)\mathrm{d}x,\ U\in\left(0,\frac{T}{\ln T}\right], \tag{4.1}$$

where $t-\varphi_1(t)\sim(1-c)\pi(t)$.

Remark 4. The formula (4.1) remains true also in the case when the integral on the right-hand side of eq. (4.1) is only relatively convergent improper integral of the second kind (in the Riemann sense).

In the case (comp. (1.7)) $T=\varphi_1(\overset{\circ}{T}),\ T+U=\varphi_1(\overset{\circ}{\widehat{T+U}})$ we obtain from (4.1)

$$\int_{\overset{\circ}{T}}^{\overset{\circ}{\widehat{T+U}}} f[\varphi_1(t)]\tilde{Z}^2(t)\mathrm{d}t=\int_T^{T+U} f(x)\mathrm{d}x. \tag{4.2}$$

4.2. First of all, we have from (4.2), for example,

$$\int_{\mathring{t}_{2\nu}(-x)}^{\mathring{t}_{2\nu}(x)} f[\varphi_1(t)]\tilde{Z}^2(t)\mathrm{d}t=\int_{t_{2\nu}(-x)}^{t_{2\nu}(x)} f(t)\mathrm{d}t, \tag{4.3}$$

(see (1.4). Next, in the case

$$f(t)=Z[\varphi_1(t)]\tilde{Z}^2(t),\ t\in \mathring{G}_1(x)\cup\mathring{G}_2(y)$$

we have the following $\tilde{Z}^2$-transformation

$$\begin{aligned}&\int_{\mathring{G}_1(x)} Z[\varphi_1(t)]\tilde{Z}^2(t)\mathrm{d}t=\int_{G_1(x)} Z(t)\mathrm{d}t, \\ &\int_{\mathring{G}_2(y)} Z[\varphi_1(t)]\tilde{Z}^2(t)\mathrm{d}t=\int_{G_2(y)} Z(t)\mathrm{d}t,\end{aligned} \tag{4.4}$$

(see (1.1), (4.3)). Let us remind that we have proved the following mean-value formulae (see [2])

$$\begin{aligned}&\int_{G_1(x)} Z(t)\mathrm{d}t=\frac{2}{\pi}H\sin x+\mathcal{O}(T^{1/6+\epsilon}), \\ &\int_{G_2(y)} Z(t)\mathrm{d}t=-\frac{2}{\pi}H\sin y+\mathcal{O}(T^{1/6+\epsilon}).\end{aligned} \tag{4.5}$$

Now, our formulae (1.5) follow from (4.4) by (4.5).

I would like to thank Michal Demetrian for helping me with the electronic version of this work.

REFERENCES

[1] A. Ivic, 'The Riemann zeta-function', A Willey-Interscience Pub., New York, 1985.

[2] J. Moser, 'New consequences of the Riemann-Siegel formula', Acta Arith., 42, (1982), 1-10 (in russian).

[3] J. Moser, 'Jacob's ladders and the almost exact asymptotic representation of the Hardy-Littlewood integral', (2008), arXiv:0901.3973.

[4] J. Moser, 'Jacob's ladders and the tangent law for short parts of the Hardy-Littlewood integral', (2009), arXiv:0906.0659.

[5] J. Moser, 'Jacob's ladders and the multiplicative asymptotic formula for short and microscopic parts of the Hardy-Littlewood integral', (2009), arXiv:0907.0301.

[6] J. Moser, 'Jacob's ladders and the quantization of the Hardy-Littlewood integral', (2009), arXiv:0909.3928.

[7] J. Moser, 'Jacob's ladders and the first asymptotic formula for the expression of the sixth order $|\zeta(1/2+i\varphi(t)/2)|^4|\zeta(1/2+it)|^2$', (2009), arXiv:0911.1246.

[8] J. Moser, 'Jacob's ladders and the first asymptotic formula for the expression of the fifth order $Z[\varphi(t)/2+\rho_1]Z[\varphi(t)/2+\rho_2]Z[\varphi(t)/2+\rho_3]\hat{Z}^2(t)$ for the collection of disconnected sets', (2009), arXiv:0912.0130.

[9] J. Moser, 'Jacob's ladders, the iterations of Jacob's ladder $\varphi_1^k(t)$ and asymptotic formulae for the integrals of the products $Z^2[\varphi_1^n(t)]Z^2[\varphi^{n-1}(t)]\cdots Z^2[\varphi_1^0(t)]$ for arbitrary fixed $n\in\mathbb{N}$' (2010), arXiv:1001.1632.

[10] J. Moser, 'Jacob's ladders and the asymptotic formula for the integral of the eight order expression $|\zeta(1/2+i\varphi_2(t))|^4|\zeta(1/2+it)|^4$', (2010), arXiv:1001.2114.

[11] J. Moser, 'Jacob's ladders and the asymptotically approximate solutions of a nonlinear diophantine equation', (2010), arXiv: 1001.3019.

[12] J. Moser, 'Jacob's ladders and the asymptotic formula for short and microscopic parts of the Hardy-Littlewood integral of the function $|\zeta(1/2+it)|^4$', (2010), arXiv:1001.4007.

[13] J. Moser, 'Jacob's ladders and the nonlocal interaction of the function $|\zeta(1/2+it)|$ with $\arg\zeta(1/2+it)$ on the distance $\sim(1-c)\pi(t)$', (2010), arXiv: 1004.0169.

[14] J. Moser, 'Jacob's ladders and the $\tilde{Z}^2$ - transformation of polynomials in $\ln\varphi_1(t)$', (2010), arXiv: 1005.2052.

[15] J. Moser, 'Jacob's ladders and the oscillations of the function $|\zeta\left(\frac{1}{2}+it\right)|^2$ around the main part of its mean-value; law of the almost exact equality of the corresponding areas', (2010), arxiv:

DEPARTMENT OF MATHEMATICAL ANALYSIS AND NUMERICAL MATHEMATICS, COMENIUS UNIVERSITY, MLYNSKA DOLINA M105, 842 48 BRATISLAVA, SLOVAKIA

E-mail address: `jan.mozer@fmph.uniba.sk`